

US009468464B2

(12) **United States Patent**
Shadduck

(10) **Patent No.:** **US 9,468,464 B2**
(45) **Date of Patent:** **Oct. 18, 2016**

(54) **METHODS FOR TREATING THE SKIN USING VACUUM**

USPC 606/131, 133, 137, 139, 172, 180, 79
See application file for complete search history.

(71) Applicant: **Axia MedSciences, LLC**, Tiburon, CA (US)

(56) **References Cited**

(72) Inventor: **John H. Shadduck**, Tiburon, CA (US)

U.S. PATENT DOCUMENTS

(73) Assignee: **Axia MedSciences, LLC**, Tiburon, CA (US)

2,608,032 A 8/1952 Garver
2,631,583 A 3/1953 Lavergne

(Continued)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

FOREIGN PATENT DOCUMENTS

DE 34 21 390 A1 12/1985
DE 234 608 4/1986

(Continued)

(21) Appl. No.: **14/702,486**

(22) Filed: **May 1, 2015**

OTHER PUBLICATIONS

(65) **Prior Publication Data**
US 2015/0230824 A1 Aug. 20, 2015

Ex Parte Reexamination Certificate U.S. Pat. No. 6,241,739 C1, Microdermabrasion Device and Method of Treating the Skin Surface, Inventor Stephen H. Waldron, Dec. 11, 2007, and file history through Aug. 8, 2006.

(Continued)

Related U.S. Application Data

(60) Continuation of application No. 13/620,164, filed on Sep. 14, 2012, which is a continuation of application No. 11/739,615, filed on Apr. 24, 2007, now Pat. No. 8,337,513, which is a division of application No.

Primary Examiner — Vy Bui

(74) *Attorney, Agent, or Firm* — Knobbe, Martens, Olson & Bear, LLP

(Continued)

(51) **Int. Cl.**
A61B 17/50 (2006.01)
A61B 17/54 (2006.01)

(Continued)

(52) **U.S. Cl.**
CPC **A61B 17/54** (2013.01); **A61B 17/545** (2013.01); **A61M 1/0064** (2013.01); **A61M 35/00** (2013.01); **A61B 2017/00761** (2013.01);


(Continued)

(58) **Field of Classification Search**
CPC A61B 17/50

(57) **ABSTRACT**

An instrument and technique for the removal of epidermal layers in a controlled manner utilizing a hand-held instrument with a working end that (i) a vacuum aspiration system, (ii) a source for delivery of a sterile fluids or pharmacological agents to the skin; and (iii) a skin interface surface in the working end that has specially shape structure for abrading surface layers of the patient's epidermis as the working end is moved over the patient's skin while at the same time causing rapid penetration of the fluids into the skin for therapeutic purposes. Movement of the working end across the skin causes abrasion of the surface layers in a path over the patient's skin.

20 Claims, 9 Drawing Sheets



Related U.S. Application Data

- 10/699,747, filed on Nov. 3, 2003, now Pat. No. 7,789,886, which is a continuation of application No. 09/648,025, filed on Aug. 25, 2000, now Pat. No. 6,641,591.
- (60) Provisional application No. 60/150,782, filed on Aug. 26, 1999.
- (51) **Int. Cl.**
A61M 1/00 (2006.01)
A61M 35/00 (2006.01)
A61B 17/32 (2006.01)
A61B 17/00 (2006.01)
- (52) **U.S. Cl.**
 CPC *A61B 2017/320004* (2013.01); *A61B 2217/005* (2013.01); *A61B 2217/007* (2013.01)

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,701,559 A	2/1955	Cooper	5,368,581 A	11/1994	Smith et al.
2,712,823 A	7/1955	Kurtin	5,391,151 A	2/1995	Wilmot
2,867,214 A	1/1959	Wilson	5,417,674 A	5/1995	Smith et al.
2,881,763 A	4/1959	Robbins	5,419,772 A	5/1995	Teitz et al.
2,921,585 A	1/1960	Schumann	5,460,620 A	10/1995	Smith et al.
3,085,573 A	4/1963	Meyer et al.	5,470,323 A	11/1995	Smith et al.
3,214,869 A	11/1965	Stryker	5,484,427 A	1/1996	Gibbons
3,476,112 A	11/1969	Elstein	5,562,642 A	10/1996	Smith et al.
3,574,239 A	4/1971	Sollerud	5,611,687 A	3/1997	Wagner
3,715,838 A	2/1973	Young et al.	5,612,797 A	3/1997	Liang et al.
3,865,352 A	2/1975	Nelson et al.	5,674,235 A	10/1997	Parisi
3,948,265 A	4/1976	Al Ani	5,676,643 A	10/1997	Cann et al.
3,964,212 A	6/1976	Karden	5,676,648 A	10/1997	Henley
3,977,084 A	8/1976	Sloan	5,683,971 A	11/1997	Rose et al.
4,121,388 A	10/1978	Wilson	5,697,920 A	12/1997	Gibbons
4,155,721 A	5/1979	Fletcher	5,707,383 A	1/1998	Bays
4,170,821 A	10/1979	Booth	5,713,785 A	2/1998	Nishio
4,182,329 A	1/1980	Smit et al.	5,735,833 A	4/1998	Olson
4,203,431 A	5/1980	Abura et al.	5,759,185 A	6/1998	Grinberg
4,216,233 A	8/1980	Stein	5,779,519 A	7/1998	Oliver
4,299,219 A	11/1981	Norris, Jr.	5,800,446 A	9/1998	Banuchi
4,378,804 A	4/1983	Cortese	5,807,353 A	9/1998	Schmitz
4,560,373 A	12/1985	Sugino et al.	5,810,842 A	9/1998	Di Fiore et al.
4,646,480 A	3/1987	Williams	5,813,416 A	9/1998	Rudolph
4,646,482 A	3/1987	Chitjian	5,817,050 A	10/1998	Klein
4,655,743 A	4/1987	Hyde	5,846,215 A	12/1998	Zygmont
4,676,749 A	6/1987	Mabille	5,848,998 A	12/1998	Marasco, Jr.
4,706,676 A	11/1987	Peck	5,861,142 A	1/1999	Schick
4,754,756 A	7/1988	Shelanski	5,873,881 A	2/1999	McEwen et al.
4,757,814 A	7/1988	Wang et al.	5,879,323 A	3/1999	Henley
4,764,362 A	8/1988	Barchas	5,882,201 A	3/1999	Salem
4,795,421 A	1/1989	Blasius, Jr. et al.	5,885,260 A	3/1999	Mehl, Sr. et al.
4,875,287 A	10/1989	Creasy et al.	5,908,401 A	6/1999	Henley
4,886,078 A	12/1989	Shiffman	5,919,152 A	7/1999	Zygmont
4,887,994 A	12/1989	Bedford	5,954,730 A	9/1999	Bernabei
4,900,316 A	2/1990	Yamamoto	5,971,999 A	10/1999	Naldoni
4,917,086 A	4/1990	Feltovich et al.	5,980,555 A	11/1999	Barbut et al.
4,925,450 A	5/1990	Imonti et al.	6,019,749 A	2/2000	Fields et al.
4,957,747 A	9/1990	Stiefel	6,024,733 A	2/2000	Eggers et al.
5,006,004 A	4/1991	Dirksing et al.	6,027,402 A	2/2000	Oliver
5,006,339 A	4/1991	Bargery et al.	6,039,745 A	3/2000	Di Fiore et al.
5,012,797 A	5/1991	Liang et al.	6,042,552 A	3/2000	Cornier
5,035,089 A	7/1991	Tillman et al.	6,080,165 A	6/2000	DeJacma
5,037,431 A	8/1991	Summers et al.	6,080,166 A	6/2000	McEwen et al.
5,037,432 A	8/1991	Molinari	6,090,085 A	7/2000	Mehl, Sr. et al.
5,100,412 A	3/1992	Rosso	6,120,512 A	9/2000	Bernabei
5,100,424 A	3/1992	Jang	6,129,701 A	10/2000	Cimino
5,119,839 A	6/1992	Rudolph	6,136,008 A	10/2000	Becker et al.
5,122,153 A	6/1992	Harrel	6,139,553 A	10/2000	Dotan
5,207,234 A	5/1993	Rosso	6,139,554 A	10/2000	Karkar et al.
5,222,956 A	6/1993	Waldron	6,142,155 A	11/2000	Rudolph
5,242,433 A	9/1993	Smith et al.	6,149,634 A	11/2000	Bernabei
5,254,109 A	10/1993	Smith et al.	6,159,226 A	12/2000	Kim
			6,162,218 A	12/2000	Elbrecht et al.
			6,162,232 A	12/2000	Shaddock
			6,183,451 B1	2/2001	Mehl, Sr. et al.
			6,183,483 B1	2/2001	Chang
			6,193,589 B1 *	2/2001	Khalaj 451/102
			6,231,593 B1	5/2001	Meserol
			6,235,039 B1	5/2001	Parkin et al.
			6,238,275 B1	5/2001	Metcalf et al.
			6,241,739 B1	6/2001	Waldron
			6,264,666 B1	7/2001	Coleman et al.
			6,277,128 B1	8/2001	Muldner
			6,283,978 B1	9/2001	Cheski et al.
			6,299,620 B1	10/2001	Shaddock
			6,306,119 B1	10/2001	Weber et al.
			6,306,147 B1	10/2001	Bernabei et al.
			6,322,568 B1	11/2001	Bernabei et al.
			6,368,333 B2	4/2002	Bernabei et al.
			6,387,103 B2	5/2002	Shaddock
			6,409,736 B1	6/2002	Bernabei
			6,410,599 B1	6/2002	Johnson
			RE37,796 E	7/2002	Henley
			6,414,032 B1	7/2002	Johnson
			6,420,431 B1	7/2002	Johnson
			6,423,078 B1	7/2002	Bays et al.
			6,423,750 B1	7/2002	Johnson
			6,432,113 B1	8/2002	Parkin et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

6,432,114 B1	8/2002	Rosso	7,678,120 B2	3/2010	Shaddock
6,471,712 B2	10/2002	Burres	7,744,582 B2	6/2010	Sadowski et al.
6,477,410 B1	11/2002	Henley et al.	7,789,886 B2	9/2010	Shaddock
6,482,212 B1	11/2002	Bernabei et al.	7,837,695 B2	11/2010	Hart et al.
6,488,646 B1	12/2002	Zygmunt	7,901,373 B2	3/2011	Tavger
6,494,856 B1	12/2002	Zygmunt	7,951,156 B2	5/2011	Karasiuk
6,500,183 B1	12/2002	Waldron	8,025,669 B1	9/2011	David et al.
6,503,256 B2	1/2003	Parkin et al.	RE42,960 E	11/2011	Waldron
6,511,486 B2	1/2003	Mercier et al.	8,048,089 B2	11/2011	Ignon et al.
6,514,262 B1	2/2003	Di Fiore et al.	8,066,716 B2	11/2011	Shaddock
6,527,783 B1	3/2003	Ignon	8,088,085 B2	1/2012	Thiebaut et al.
6,535,761 B2	3/2003	Bernabei	8,128,638 B2	3/2012	Karasiuk et al.
6,540,757 B1	4/2003	Hruska et al.	8,221,437 B2	7/2012	Waldron et al.
6,562,013 B1	5/2003	Marasco, Jr.	8,236,008 B2	8/2012	Boone, III et al.
6,562,050 B1	5/2003	Owen	8,277,287 B2	10/2012	Hart
6,564,093 B1	5/2003	Tannenbaum et al.	8,337,513 B2	12/2012	Shaddock
6,565,535 B2	5/2003	Zaias et al.	8,343,116 B2	1/2013	Ignon et al.
6,582,442 B2	6/2003	Simon et al.	8,814,836 B2	8/2014	Ignon et al.
6,589,218 B2	7/2003	Garcia	9,056,193 B2	6/2015	Ignon et al.
6,592,595 B1	7/2003	Ignon et al.	2001/0023351 A1	9/2001	Eilers
6,629,983 B1	10/2003	Ignon	2001/0037118 A1	11/2001	Shaddock
6,641,591 B1	11/2003	Shaddock	2001/0049511 A1	12/2001	Coleman et al.
6,645,184 B1	11/2003	Zelickson et al.	2002/0016601 A1	2/2002	Shaddock
6,652,888 B2	11/2003	Rhoades	2002/0041891 A1	4/2002	Cheski
6,673,081 B1	1/2004	Tavger et al.	2002/0058952 A1	5/2002	Weber et al.
6,673,082 B2	1/2004	Mallett et al.	2002/0107527 A1	8/2002	Burres
6,685,853 B1	2/2004	Angelopoulos et al.	2002/0133110 A1	9/2002	Citow
6,687,537 B2	2/2004	Bernabei	2002/0133176 A1	9/2002	Parkin et al.
6,695,853 B2	2/2004	Karasiuk	2002/0151826 A1	10/2002	Ramey et al.
6,735,470 B2	5/2004	Henley et al.	2002/0151908 A1	10/2002	Mallett, Sr. et al.
6,743,211 B1	6/2004	Prausnitz et al.	2002/0188261 A1	12/2002	Hruska
6,743,215 B2	6/2004	Bernabei	2003/0012415 A1	1/2003	Cossel
6,764,493 B1	7/2004	Weber et al.	2003/0018252 A1	1/2003	Duchon et al.
6,869,611 B1	3/2005	Kligman et al.	2003/0060834 A1	3/2003	Muldner
6,905,487 B2	6/2005	Zimmerman	2003/0093040 A1	5/2003	Miksza et al.
6,911,031 B2	6/2005	Muldner	2003/0093089 A1	5/2003	Greenberg
6,924,649 B2	8/2005	Knoedgen	2003/0097139 A1	5/2003	Karasiuk
6,926,681 B1	8/2005	Ramey et al.	2003/0167032 A1	9/2003	Ignon
6,942,649 B2	9/2005	Ignon et al.	2003/0187462 A1	10/2003	Chang
7,001,355 B2	2/2006	Nunomura et al.	2003/0208159 A1	11/2003	Ignon et al.
7,004,933 B2	2/2006	McDaniel	2003/0212127 A1	11/2003	Glassman et al.
7,044,938 B2	5/2006	La Bianco et al.	2003/0212415 A1	11/2003	Karasiuk
7,052,503 B2	5/2006	Bernabei	2004/0010222 A1	1/2004	Nunomura et al.
7,069,073 B2	6/2006	Henley et al.	2004/0010269 A1	1/2004	Grimes et al.
7,070,488 B2	7/2006	Suissa et al.	2004/0087972 A1	5/2004	Mulholland et al.
7,083,580 B2	8/2006	Bernabei	2004/0092895 A1	5/2004	Harmon
7,087,063 B2	8/2006	Carson et al.	2004/0092959 A1	5/2004	Bernaz
7,094,252 B2	8/2006	Koop	2004/0097967 A1	5/2004	Ignon
7,115,275 B2	10/2006	Clarot et al.	2004/0122447 A1	6/2004	Harmon et al.
7,135,011 B2	11/2006	Powers et al.	2004/0143274 A1	7/2004	Shaddock
7,153,311 B2	12/2006	Chung	2004/0162565 A1	8/2004	Carson et al.
7,197,359 B1	3/2007	Tokudome et al.	2004/0166172 A1	8/2004	Rosati et al.
7,198,623 B2	4/2007	Fischer et al.	2004/0219179 A1	11/2004	McDaniel
7,232,444 B2	6/2007	Chang	2004/0236291 A1	11/2004	Zelickson et al.
7,241,208 B2	7/2007	Suissa et al.	2004/0243149 A1	12/2004	Lee, Jr.
7,276,051 B1	10/2007	Henley et al.	2004/0254587 A1	12/2004	Park
7,314,326 B2	1/2008	Rosenberg	2004/0267285 A1	12/2004	Chang
7,316,657 B2	1/2008	Kleinhenz et al.	2005/0037034 A1	2/2005	Rhoades
7,318,828 B1	1/2008	Revivo	2005/0038448 A1	2/2005	Chung
7,320,691 B2	1/2008	Pilcher et al.	2005/0059940 A1	3/2005	Weber et al.
7,320,801 B2	1/2008	Kelly	2005/0084509 A1	4/2005	Bernstein
7,354,423 B2	4/2008	Zelickson et al.	2005/0148958 A1	7/2005	Rucinski
7,364,565 B2	4/2008	Freeman	2005/0203111 A1	9/2005	David
7,384,405 B2	6/2008	Rhoades	2005/0209611 A1	9/2005	Greenberg
7,427,273 B2	9/2008	Mitsui	2005/0283176 A1	12/2005	Law
7,458,944 B2	12/2008	Liste et al.	2006/0002960 A1	1/2006	Zoeteweij et al.
7,476,205 B2	1/2009	Erdmann	2006/0116674 A1	6/2006	Goble et al.
7,477,938 B2	1/2009	Sun et al.	2006/0161178 A1	7/2006	Lee
7,482,314 B2	1/2009	Grimes et al.	2006/0189964 A1	8/2006	Anderson
7,489,989 B2	2/2009	Sukhanov et al.	2006/0191562 A1	8/2006	Numomura
7,507,228 B2	3/2009	Sun et al.	2006/0200099 A1	9/2006	La Bianco et al.
7,582,067 B2	9/2009	Van Acker	2006/0200172 A1	9/2006	Shaddock
7,597,900 B2	10/2009	Zimmer et al.	2006/0200173 A1	9/2006	Shaddock
7,597,901 B2	10/2009	Clarot et al.	2006/0212029 A1	9/2006	Arcusa Villacampa et al.
7,658,742 B2	2/2010	Karasiuk	2006/0253125 A1	11/2006	Ignon
			2006/0264893 A1	11/2006	Sage, Jr. et al.
			2007/0005078 A1	1/2007	Hart et al.
			2007/0043382 A1	2/2007	Cheney
			2007/0065515 A1	3/2007	Key

(56)

References Cited

U.S. PATENT DOCUMENTS

2007/0088371 A1 4/2007 Karasiuk
 2007/0123808 A1 5/2007 Rhoades
 2007/0154502 A1 7/2007 Hattendorf et al.
 2007/0156124 A1 7/2007 Ignon et al.
 2007/0178121 A1 8/2007 First et al.
 2007/0198031 A1 8/2007 Kellogg
 2007/0208353 A1 9/2007 Shadduck
 2007/0239173 A1 10/2007 Khalaj
 2008/0027328 A1 1/2008 Klopotek et al.
 2008/0091179 A1 4/2008 Durkin et al.
 2008/0103563 A1 5/2008 Powell
 2008/0119781 A1 5/2008 King
 2008/0132914 A1 6/2008 Bossard et al.
 2008/0139974 A1 6/2008 Da Silva
 2008/0154161 A1 6/2008 Abbott
 2008/0193493 A1 8/2008 Rhoades
 2008/0200861 A1 8/2008 Shalev et al.
 2008/0208146 A1 8/2008 Brandwein et al.
 2008/0214987 A1 9/2008 Xu
 2008/0215068 A1 9/2008 Hart et al.
 2008/0221548 A1 9/2008 Danenberg et al.
 2008/0243039 A1 10/2008 Rhoades
 2008/0287864 A1 11/2008 Rosenberg
 2008/0300529 A1 12/2008 Reinstein
 2008/0300552 A1 12/2008 Cichocki et al.
 2009/0048557 A1 2/2009 Yeshurun et al.
 2009/0053390 A1 2/2009 Sakou et al.
 2009/0062815 A1 3/2009 Karasiuk et al.
 2009/0099091 A1 4/2009 Hantash
 2009/0099093 A1 4/2009 Hantash
 2009/0124985 A1 5/2009 Hasenoehrl et al.
 2009/0138026 A1 5/2009 Wu
 2009/0177171 A1 7/2009 Ignon et al.
 2009/0192442 A1 7/2009 Ignon et al.
 2009/0222023 A1 9/2009 Boone, III et al.
 2010/0045427 A1 2/2010 Boone, III et al.
 2010/0049177 A1 2/2010 Boone, III et al.
 2010/0049210 A1 2/2010 Boone, III et al.
 2010/0217357 A1 8/2010 Da Silva
 2011/0054490 A1 3/2011 Hart
 2011/0066162 A1 3/2011 Cohen
 2011/0082415 A1 4/2011 Ignon et al.
 2012/0022435 A1 1/2012 Ignon et al.
 2012/0041338 A1 2/2012 Chickering, III et al.
 2012/0136374 A1 5/2012 Karasiuk
 2013/0018317 A1 1/2013 Bobroff et al.
 2013/0066336 A1 3/2013 Boone, III et al.
 2013/0096577 A1 4/2013 Shadduck
 2013/0102978 A1 4/2013 Ignon et al.
 2013/0144280 A1 6/2013 Eckhouse et al.
 2013/0158547 A1 6/2013 David
 2014/0343481 A1 11/2014 Ignon
 2014/0343574 A1 11/2014 Ignon et al.
 2015/0032047 A1 1/2015 Ignon et al.

2015/0230824 A1 8/2015 Shadduck
 2015/0230825 A1 8/2015 Shadduck
 2015/0231379 A1 8/2015 Ignon et al.
 2015/0265822 A1 9/2015 Ignon et al.
 2015/0272623 A1 10/2015 Ignon et al.
 2015/0290442 A1 10/2015 Ignon et al.
 2016/0038183 A1 2/2016 Ignon et al.

FOREIGN PATENT DOCUMENTS

DE 10 2004 015815 A1 11/2005
 EP 0 258 901 9/1987
 EP 0 564 392 3/1993
 IT 553 076 12/1956
 IT 118 49 22 3/1985
 JP 1993-088552 12/1993
 JP 09-294747 11/1997
 JP 2003-339713 12/2003
 JP 2006-503627 2/2006
 JP 2006-204767 10/2006
 KR 20-0280320 7/2002
 WO WO 00/15300 3/2000
 WO WO 01/93931 12/2001
 WO WO 03/073917 9/2003
 WO WO 2004/037098 5/2004
 WO WO 2005/070313 8/2005
 WO WO 2006/018731 2/2006
 WO WO 2007/114904 10/2007
 WO WO 2012/145667 10/2012

OTHER PUBLICATIONS

File History of Reissue No. 11/027,590, filed Dec. 29, 2004 (Reissue of U.S. Pat. No. 6,500,183, issued Dec. 31, 2002).
 File History of Reexamination No. 90/007,683 (Reexamination of U.S. Pat. No. 6,241,739, issued Jun. 5, 2001).
 Cox III et al., *Decreased Splatter in Dermabrasion*, Arch Facial Plastic Surgery, Jan.-Mar. 2000, vol. 2, pp. 23-26.
 Ditre et al., *Effect of α -hydroxy acids on photoaged skin: A pilot clinical, histologic, and ultrastructural study*, Journal of American Academy of Dermatology, Feb. 1996, vol. 34, No. 2, Part 1, pp. 187-195.
 Harris et al., *Combining Manual Dermabrasion with Low Strength Trichloroacetic Acid to Improve Antinically Injured Skin*, The Journal of Dermatologic Surgery and Oncology, Jul. 1994, vol. 20, No. 7, pp. 436-442.
 File History of Reissue U.S. Appl. No. 11/027,590, filed Dec. 29, 2004 (Reissue of U.S. Pat. No. 6,500,183, issued Dec. 31, 2002).
 File History of Reexamination U.S. Appl. No. 90/007,683, filed Aug. 22, 2005 (Reexamination of U.S. Pat. No. 6,241,739, issued Jun. 5, 2001).
 File History of Reexamination U.S. Appl. No. 90/013,284, filed Jul. 2, 2014 (Reexamination of U.S. Pat. No. 6,241,739, issued Jun. 5, 2001).

* cited by examiner




FIG. 1A




FIG. 1B




FIG. 2




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9

METHODS FOR TREATING THE SKIN USING VACUUM

PRIORITY INFORMATION

This application is a continuation of U.S. patent application Ser. No. 13/620,164, filed Sep. 14, 2012, which is a continuation of U.S. patent application Ser. No. 11/739,615, filed Apr. 24, 2007, now U.S. Pat. No. 8,337,513, which is a divisional of U.S. patent application Ser. No. 10/699,747, filed Nov. 3, 2003, now U.S. Pat. No. 7,789,886, which is a continuation of U.S. patent application Ser. No. 09/648,025 filed Aug. 25, 2000, now U.S. Pat. No. 6,641,591, which claims the priority benefit under 35 U.S.C. §119(e) of Provisional U.S. Patent Application No. 60/150,782, filed Aug. 26, 1999, the entire contents of these applications being hereby incorporated by reference herein.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to devices for dermatology and more particularly to a hand-held instrument with a working end that carries (i) a negative pressure aspiration system, (ii) a source for delivery of a sterile fluids to the skin; and (iii) a skin interface surface in the working end that has specially shape structure for abrading surface layers of the patient's epidermis as the working end is moved over the patient's skin while at the same time causing rapid penetration of the fluids into the skin for therapeutic purposes.

2. Description of the Related Art

Dermatologists and plastic surgeons have used various methods for removing superficial skin layers to cause the growth of new skin layers (i.e., commonly described as skin resurfacing techniques) since the early 1900's. Early skin resurfacing treatments used an acid such as phenol to etch away surface layers of a patient's skin that contained damage to thereafter be replaced by new skin. (The term damage when referring to a skin disorder is herein defined as any cutaneous defect, e.g., including but not limited to rhytides, hyperpigmentation, acne scars, solar elastosis, other dyschromias, stria distensae, seborrheic dermatitis).

Following the removal of surface skin layers at a particular depth, no matter the method of skin removal, the body's natural wound-healing response begins to regenerate the epidermis and underlying wounded skin layers. The new skin layer will then cytologically and architecturally resemble a younger and more normal skin. The range of resurfacing treatments can be divided generally into three categories based on the depth of the skin removal and wound: (i) superficial exfoliations or peels extending into the epidermis, (ii) medium-depth resurfacing treatments extending into the papillary dermis, and (iii) deep resurfacing treatments that remove tissue to the depth of the reticular dermis (see FIGS. 1A-1B).

Modern techniques for skin layer removal include: CO.sub.2 laser resurfacing which falls into the category of a deep resurfacing treatment; Erbium laser resurfacing which generally is considered a medium-depth treatment; mechanical dermabrasion using high-speed abrasive wheels which results in a medium-depth or deep resurfacing treatment; and chemical peels which may range from a superficial to a deep resurfacing treatment, depending on the treatment parameters. A recent treatment, generally called micro-dermabrasion, has been developed that uses an air-pressure source to deliver abrasive particles directly against a patient's skin at high-velocities to abrade away skin layers.

Such a micro-dermabrasion modality may be likened to sandblasting albeit at velocities that do not cause excess pain and discomfort to the patient. Micro-dermabrasion as currently practiced falls into the category of a superficial resurfacing treatment.

A superficial exfoliation, peel or abrasion removes some or all of the epidermis (see FIGS. 1A-1B) and thus is suited for treating very light rhytides. Such a superficial exfoliation is not effective in treating many forms of damage to skin. A medium-depth resurfacing treatment that extends into the papillary dermis (see FIG. 1B) can treat many types of damage to skin. Deep resurfacing treatments, such as CO.sub.2 laser treatments, that extend well into the reticular dermis (see FIG. 1B) causes the most significant growth of new skin layers but carry the risk of scarring unless carefully controlled.

It is useful to briefly explain the body's mechanism of actually resurfacing skin in response to the removal of a significant depth of dermal layers. Each of the above-listed depths of treatment disrupts the epidermal barrier, or a deeper dermal barrier (papillary or reticular), which initiates varied levels of the body's wound-healing response. A superficial skin layer removal typically causes a limited wound-healing response, including a transient inflammatory response and limited collagen synthesis within the dermis. In a medium-depth or a deep treatment, the initial inflammatory stage leads to hemostasis through an activated coagulation cascade. Chemotactic factors and fibrin lysis products cause neutrophils and monocytes to appear at the site of the wound. The neutrophils sterilize the wound site and the monocytes convert to macrophages and elaborate growth factors which initiate the next phase of the body's wound-healing response involving granular tissue formation. In this phase, fibroblasts generate a new extracellular matrix, particularly in the papillary and reticular dermis, which is sustained by angiogenesis and protected anteriorly by the reforming epithelial layer. The new extracellular matrix is largely composed of collagen fibers (particularly Types I and III) which are laid down in compact parallel arrays (see FIG. 1B). It is largely the collagen fibers that provide the structural integrity of the new skin—and contribute to the appearance of youthful skin.

All of the prevalent types of skin damage (rhytides, solar elastosis effects, hyperpigmentation, acne scars, dyschromias, melasma, stria distensae) manifest common histologic and ultrastructural characteristics, which in particular include disorganized and thinner collagen aggregates, abnormalities in elastic fibers, and abnormal fibroblasts, melanocytes and keratinocytes that disrupt the normal architecture of the dermal layers. It is well recognized that there will be a clinical improvement in the condition and appearance of a patient's skin when a more normal architecture is regenerated by the body's wound-healing response. Of most significance to a clinical improvement is skin is the creation of more dense parallel collagen aggregates with decreased periodicity (spacing between fibrils). The body's wound-healing response is responsible for synthesis of these collagen aggregates. In addition to the body's natural wound healing response, adjunct pharmaceutical treatments that are administered concurrent with, or following, a skin exfoliations can enhance the development of collagen aggregates to provide a more normal dermal architecture in the skin—the result being a more youthful appearing skin.

The deeper skin resurfacing treatments, such as laser ablation, chemical peels and mechanical dermabrasion have drawbacks. The treatments are best used for treatments of a patient's face and may not be suited for treating other

portions of a patient's body. For example, laser resurfacing of a patient's neck or décolletage may result in post-treatment pigmentation disorders. All the deep resurfacing treatments are expensive, require anesthetics, and must be performed in a clinical setting. Perhaps, the most significant disadvantage to deep resurfacing treatments relates to the post-treatment recovery period. It may require up to several weeks or even months to fully recover and to allow the skin the form a new epidermal layer. During a period ranging from a few weeks to several weeks after a deep resurfacing treatment, the patient typically must wear heavy make-up to cover redness thus making the treatment acceptable only to women.

The superficial treatment offered by micro-dermabrasion has the advantages of being performed without anesthetics and requiring no extended post-treatment recovery period. However, micro-dermabrasion as currently practices also has several disadvantages. First, a micro-dermabrasion treatment is adapted only for a superficial exfoliation of a patient's epidermis which does not treat many forms of damage to skin. Further, the current micro-dermabrasion devices cause abrasive effects in a focused area of the skin that is very small, for example a few mm.^{sup.2}, since all current devices use a single pin-hole orifice that jets air and abrasives to strike the skin in a highly focused area. Such a focused treatment area is suitable for superficial exfoliations when the working end of the device is passed over the skin in overlapping paths. Further, such focused energy delivery is not well suited for deeper skin removal where repeated passes may be necessary. Still further, current micro-dermabrasion devices are not suited for deeper skin removal due to the pain associated with deep abrasions. Other disadvantages of the current micro-dermabrasion devices relate to the aluminum oxide abrasive particles that are typically used. Aluminum oxide can contaminate the working environment and create a health hazard for operators and patients alike. Inhalation of aluminum oxide particles over time can result in serious respiratory disorders.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B are sectional illustrations of a patient's skin showing dermal layers.

FIG. 2 is a view of a Type "A" body and working end of the instrument of the invention.

FIG. 3 is an enlarged view of the working end of the instrument of FIG. 2.

FIG. 4 is a sectional view of working end of FIG. 3.

FIG. 5 is a view showing the manner of using the working end of the invention of FIGS. 3-4 in performing a method of the invention.

FIG. 6 is a view of a Type "B" body, working end and handle in an exploded view.

FIG. 7 is a view of the working end of the instrument of FIG. 6 and a housing.

FIG. 8 is an enlarged view of the skin interface of the working end of FIG. 7.

FIG. 9 is a sectional view of the skin interface of FIG. 8.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

1. Type "A" Skin Resurfacing System. Referring to FIGS. 2-3, an exemplary instrument system 5 is shown for removing superficial skin layers. The instrument system 5 includes: (i) a hand-held body 18 with a working end 20 that defines a skin interface surface portion indicated at 25 in

FIGS. 2-3. An opening portion 26 transitions into an interior passageway 28 that extends through the body to communicate with a negative (-) pressure source (or aspiration source) indicated at 30 that operates as vacuum means for aspirating skin debris from a targeted skin surface treatment site TS.

Of particular interest, FIGS. 3-4 show views of the working end 20 with the skin interface 25 being configured with a particular irregular or ridged surface structure indicated at 32. The ridged surface structure 32 further has a particular minimum width dimension W to accommodate from the ridge shape with as many as about 25 ridges on each side of opening 26 depending on the overall dimensions of the working end 20. More particular aspects of the irregular or ridged surface structure 32 will be described below.

In this preferred embodiment, the working end 20 is of any suitable material, such as a transparent medical grade plastic. The transparency of the working end will assist the operator in localizing treatment in a particular targeted skin treatment area. The overall transverse dimension of the working end 20 of FIGS. 2-3 may be from around about 5.0 mm. to about 50.0 mm. with a larger dimensioned end being adapted for treating a larger skin area (e.g., arms, back legs and décolletage). A typical dimension is from about 5.0 mm. to 15.0 mm. for a skin treatment site area TS around a patient's face.

The invention allows the area (e.g., in mm²) of opening 26 be in any selected shape but preferably is an elongate shape in the center of the working end 25. The open distal end 26 comprises the distal termination of passageway 28 and the proximal end of the passageway in handle 18 is connected to a flexible aspiration tube 33 that extends to a remote collection reservoir 35 intermediate to the actual aspiration source 30. The aspiration source 30 thus is adapted to draw the working end 20 and more particularly the skin interface 25 against the skin treatment site TS to perform the method of the invention as will be described below. The aspiration source or negative (-) pressurization source 30 may be any suitable vacuum source known in the art. Between the aspiration source 30 and remote collection reservoir 35 may be a filter 38 subsystem that is known in the art for collecting aspirated skin detritus and spent crystalline agents CA that are captured in the open distal end of passageway chamber 28. The collection reservoir 35 and filter 38 are preferably of inexpensive plastic and other materials that are disposable.

The aspiration source 30 may be provided with an adjustable valve means 40 for adjusting the pressure level setting to any suitable range. The physician will learn from experience how to balance the pressure level to attain the desired level of suction against the patient's skin. A trigger or switch component 42 is provided as a foot-switch (FIG. 2) but any suitable finger switch in the body 18 also may be used.

The working end 20 also carries means for introducing abrasive crystals into the working end or distalmost end of passageway 28 to allow individual loose crystalline agents CA to thereafter be captured between the skin interface 25 and the patient's skin. In this embodiment, two channels 44a-44b are provided together with flexible tubes 46a-46b to introduce the loose crystalline agents CA into the working end (see FIGS. 2-4). Each distal portion 47 (collectively) of the channels 44a-44b may comprise a small dimension aperture to limit the rate of flow of crystalline agents CA into the working end. The number of such channels (i.e., 44a-44n) may range from one to about ten and fall within the scope of the invention. Any singular or plural number of channels can serve the purpose of slowly introducing crystal

into the working end. Referring to FIGS. 2-3, the crystalline agent CA delivery source 50 comprises a reservoir 55 that holds a suitable volume of abrasive crystals for a single treatment or a number of treatments. A flexible supply tube 56 extends between a remote the reservoir 55, and in this embodiment the tube is split to connect to the two channels 44a-44b. Preferably, the remote reservoir 55 that carries the crystalline agent CA is unpressurized but carries air intake relief valve 58 such that any slight negative pressure created by the aspiration source 30 when the skin interface is in contact with a patient's skin will draw crystals to the working end. It should be appreciated that reservoir 55 may be built into handle body 18 and fall within the scope of the invention. The crystal delivery source 50 may carry crystals ranging in size from about 1 μm to about 50 μm in maximum cross-sectional dimension, (for example, aluminum oxide crystals). Preferably, the crystals are from about 5 μm to about 30 μm in maximum cross-sectional dimension to allow a very fine abrasion of the epidermis.

It has been found that by a slight negative pressure environment the open end 26 and passageway 28, the crystalline agent will be caused to dribble into, or be sucked into, the passageway 28 in the working end 20. Thereafter, the movement of the working end 20 in a sideways movement over the skin causes a portion of the crystalline agent CA volume to be captured temporarily in the irregular or corrugated surface structure of the skin interface 25. In this process of moving the skin interface 25 over the targeted treatment site TS, it has been found that the sharp-edged crystalline agents are rolled over and over while being pressed into the surface of the skin and thereby abrade and remove the skin surface in a controllably gentle manner that is below any threshold of significant pain.

After the spent crystals are rolled over and over by the skin interface when moving in a first lateral direction across the skin, and after the working end is then is reversed in directional movement across the skin, a portion of the spent crystals and abraded skin debris necessarily roll into the central opening portion 26 wherein the negative pressure environment captures and aspirates the abraded materials to the remote collection reservoir 35.

To facilitate the process described above, the invention is provided with novel aspects that relate to the irregular or ridged surface structure 32 mentioned above. The entire skin interface 25 may be of any suitable plan form (e.g., round, oval, rectangular etc.) and fall within the scope of the invention. More in particular, the interface 25 defines a 1st outer periphery 25A and a 2nd inner periphery 25B that generally are in apposition to one another and are spaced apart by width W with the inner periphery about the edge of opening 26 (see FIG. 3).

In a preferred embodiment shown in FIGS. 3-4, the concept of 1st and 2nd peripheries 25A and 25B in apposition thus comprise peripheries that are dual and side-by-side as shown in FIG. 4 and are thus adapted for side-to-side lateral or sideways movement while performing the technique of the invention, for example which is a natural movement of a human hand over a patient's skin. Thus, the direction of the ridges 60 extend generally transverse relative to a line drawn that indicates the direction of movement of the working end 20 in performing the method of the invention. That is, in the exemplary working end of FIG. 4, the working end is generally optimized for side-to-side or lateral movement. Thus, the ridge alignment is generally transverse to the direction of movement in operations indicated by arrow A. (In a circular working end that is adapted generally for movement in any direction, the direction of the ridges 60

may be generally transverse to any direction of movement by being concentric relative to a central opening 26 (not shown)).

The terms irregular or ridged shape structure 32 as used herein mean that a series of at least one projecting edge portion 62a projects distally as a ridge within the skin interface portion 25. The irregular shape structure 32 further typically carries recessed portions or valley portions 62b that are recessed in the proximal direction intermediate to any plurality of projecting edge portions 62a. These surface configurations for convenience are herein termed the primary shape structure (or ridge and valley elements). The width of the skin interface 25 containing shape structure 32 may be from about 2.0 mm. to 25.0 mm. or more and preferably is from about 3.0 mm. to 10.0 mm. The number of ridges preferably are from about 1 ridge to 25 ridges on each side of the opening 26. The height H of any ridge from the apex of the projecting portion 62a to the depth of the valley portion 62b may be from about 0.25 mm. to about 5.0 mm. and is preferably from about 0.5 mm. to about 2.0 mm. It has been found that various ridge height dimensions are optimal depending on the patient's skin type. Further, but optionally, it has been found that secondary shape structure of notches or recessed grooves 66 configured across the primary shape structure of ridge and valley elements may help introduce loose crystals to regions of the skin interface 25 in contact with the skin which is desirable. Such secondary grooves 66 are shown in FIG. 4 and are preferably somewhat in alignment with an axis of channels 44a-44b that introduce crystals into the working end 20 thus allowing the crystals to be suctioned into the valleys 62b of the primary shape structure.

While the series of primary ridge and valley elements together the secondary grooves seems to be optimal for the method described below, it should be appreciated that the method also may be performed with a skin interface that has (i) only primary ridge and valley elements; (ii) or only a particular surface roughness that is appropriate for partially capturing loose crystals as will be described below—as long as the skin interface has a minimum width of about 3.0 mm. which was described as a preferred width dimension previously.

FIG. 4 further shows that at least some of the crests or apexes of some of the ridge portions 62a together with the outermost periphery of the skin interface 25 define an overall tissue-receiving shape 64 that may range from flat to concave and is shown in a preferred concave configuration. The alternative shapes 64a-44b are intended to indicate an approximate range of shapes that are suitable. The apexes of ridges 62a need not all be at the same height to define shape 64. The purpose of the concave shape is to cause the outer periphery of the working end to be in firm contact with the tissue surface while the negative pressure from aspiration source 30 draws the skin into firm contact with tissue interface 25.

2. Practice of the Method of the Invention. Now turning to FIG. 5, a sectional view of working end 20 shows the technique of the present invention in exfoliating or removing skin surface layers. FIG. 5 shows the working end 20 after actuation of the negative (-) pressure source 30 with the skin surface 70 initially being drawn into the concave shape 64. The operating negative pressures may be in any suitable range that is determined by investigation. It has been found by experimentation that optimal pressure levels vary greatly depending on (i) the type of skin targeted for treatment, (ii) the dimensions across the working end, and (iii) the dimensions of opening 26.

Next, the operator moves the skin interface **25** across a treatment site TS which is a path on the patient's skin while still actuating moves the trigger **42** thereby maintaining the negative pressure environment in the passageway **26**. The negative pressure environment within the working end 5 causes crystalline particles and entrained in air to be drawn into passageway **28** proximate to the skin surface and into the shape structure **32** of the skin interface **25**. The sideways or lateral movement of the skin interface **25** captures a portion of the crystals between the interface and the skin surface, in part by over-rolling them. The continued rolling of the sharp-edged crystals trapped between the instrument and the skin surface **70** causes an abrasion and removal of the skin surface in a controllable manner.

As working end is moved in a reverse direction, the negative pressure environment in the passageway **28** captures and aspirates the spent crystals and skin debris to the remote collection reservoir **35**. At the end of a particular lateral movement of the working end, the operator may release the trigger **42** which terminates the crystal agent delivery and further allows the operator to easily lift the working end from the patient's skin. The treated path can be easily seen and the operator then can exfoliate another slightly overlapping or adjacent path by repeating the above steps until surface removal is completed over the targeted treatment area.

3. Type "B" Skin Resurfacing System. Referring to FIGS. **6-9**, another exemplary instrument system and treatment device **205** is shown for removing superficial skin layers. This system differs greatly from the Type "A" embodiment in the mechanism of action that abrades the skin since the Type "B" system uses a fluid media plus an abrading structure on the skin interface. Still several features of the Type "B" embodiment are similar to the Type "A" embodiment and the two modalities of treatment may be used to complement one another.

FIG. **6** shows that a hand-held instrument **208** has a removable working end **220** that defines a skin interface surface portion indicated at **225**. Handle portion **227a** mates with housing **227b**. A flexible tube **228** extends to a vacuum source **230**. A fluid reservoir **235** carrying a fluid skin treatment media is housed in the handle although it could also be a remote reservoir.

Referring now to FIGS. **7-9**, a first aperture arrangement consisting of at least one port or opening portion **240** of skin interface **225** that communicates with an interior passageway **242** that extends through housing **227b** to hose **228** and the vacuum or negative (-) pressure source.

FIGS. **7-9** further show a second aperture arrangement in the skin interface consisting of at least one port or openings **250** that extend around an outer periphery of the skin interface **225**. These opening(s) of the second aperture arrangement are in fluid communication with the reservoir **235** and the treatment media therein. The skin interface has a series of primary ridge elements **255a** and valley elements **255b** together the secondary notches or grooves **260** as defined above with similar dimensional parameters. This embodiment differs however in that the apexes of ridge elements **255a** are substantially a sharp edge as are the edged of the notches **260**. Thus, these primary surface elements **255a** and secondary surface elements thereby define teeth therebetween that seem well suited to abrading skin layers particularly after being hydrated by the fluid source of the system. Experimentation has shown that the vacuum source and fluid source may be reversed between the first and second aperture arrangements **240** and **250** with the method of skin removal still working well. The vacuum system

aspirates away skin debris and spent fluids as described previously. Of particular interest, the method of the invention appears to work well because the suction on the skin treatment site very quickly hydrated, or puffs up, the skin which in turn make the surface layer susceptible to painless abrasion. The ability of the system to rapidly deliver fluids to subsurface tissues allows the use of any pharmacological agent known in the art for enhancing skin rejuvenation as a part of the skin treatment. The system can use sterile water or saline solution for a treatment to remove dermal tissue with the abrasive surface of the treatment device. The system can also use a fluid carrying a chemical agent of a suitable concentration be selected from a group of acids including TCA (trichloroacetic acid), a glycolic acid including an alphahydroxy acid (AHA), a lactic acid, a citric acid, or phenol as disclosed in co-pending U.S. patent application Ser. No. 09/524,731 filed Mar. 14, 2000 which is incorporated herein by this reference.

Specific features of the invention may be shown in some figures and not in others, and this is for convenience only and any feature may be combined with another in accordance with the invention. While the principles of the invention have been made clear in the exemplary embodiments, it will be obvious to those skilled in the art that modifications of the structure, arrangement, proportions, elements, and materials may be utilized in the practice of the invention, and otherwise, which are particularly adapted to specific environments and operative requirements without departing from the principles of the invention. The appended claims are intended to cover and embrace any and all such modifications, with the limits only of the true purview, spirit and scope of the invention.

What is claimed is:

1. A method of treating a skin surface of a subject, comprising:
 - positioning a handheld device against a skin surface of the subject, the handheld device comprising:
 - a body comprising a housing;
 - a working end portion positioned along a first end of the body, the working end portion comprising a distal end configured to contact the skin surface, wherein the working end portion comprises a perimeter along the distal end;
 - a first aperture arrangement comprising at least one first port located along or near the working end portion, the at least one first port being in fluid communication with a vacuum source via at least one passageway; and
 - a second aperture arrangement comprising at least one second port located along or near the working end portion, the at least one second port being in fluid communication with a treatment media source;
 - activating the vacuum source to simultaneously deliver a treatment media to the skin surface through the at least second port, wherein the treatment media comprises a liquid, and to aspirate spent treatment media through the at least one first port and the at least one passageway, thereby selectively providing a volume of the treatment media to the skin surface of the subject;
 - wherein activating the vacuum source facilitates delivery of treatment media to subsurface skin tissue of the subject and causes adjacent skin surface to at least partially puff up to facilitate the treatment method.
2. The method of claim **1**, wherein, when in use, the vacuum source draws skin through an aperture defined by the perimeter of the working end portion to facilitate contact between the working end portion and the skin surface.

9

3. The method of claim 1, wherein the at least one passageway comprises a conduit that extends proximally from the housing of the device to the vacuum source.

4. The method of claim 1, wherein the at least one second port is in fluid communication with the treatment media source via a treatment media passageway.

5. The method of claim 1, wherein the treatment media source is separate from the handheld device.

6. The method of claim 1, wherein the treatment media source comprises a reservoir.

7. The method of claim 6, further comprising selecting a reservoir with a desired treatment media and placing said reservoir in fluid communication with the at least one second port of the handheld device.

8. The method of claim 1, wherein the treatment media comprises at least one of water and saline.

9. The method of claim 1, wherein the treatment media comprises a chemical agent.

10. The method of claim 9, wherein the chemical agent comprises at least one acid.

11. The method of claim 10, wherein the at least one acid comprises at least one of trichloroacetic acid (TCA), glycolic acid, alphahydroxy acid (AHA), lactic acid, citric acid and phenol.

12. A method of treating a skin surface of a subject, comprising:

positioning a handheld device against a skin surface of the subject, the handheld device comprising a working end portion positioned along a distal end of the handheld device, the working end portion configured to contact the skin surface, wherein the working end portion comprises an outer perimeter configured to contact the skin surface;

wherein the handheld device further comprises a at least one first port and at least one second port located along or near the working end portion, the at least one first

10

port being in fluid communication with a vacuum source, and the at least one second port being in fluid communication with a treatment media source;

activating the vacuum source to simultaneously deliver a treatment media to the skin surface through the at least second port, and to aspirate spent treatment media through the at least one first port, thereby selectively providing a volume of the treatment media to the skin surface of the subject, wherein the treatment media comprises a liquid;

wherein activating the vacuum source facilitates delivery of treatment media to subsurface skin tissue of the subject to facilitate the treatment method.

13. The method of claim 12, wherein, when in use, the vacuum source draws skin through an aperture defined by the perimeter of the working end portion to facilitate contact between the working end portion and the skin surface.

14. The method of claim 12, wherein the treatment media source is separate from the handheld device.

15. The method of claim 12, wherein the treatment media source comprises a reservoir.

16. The method of claim 15, further comprising selecting a reservoir with a desired treatment media and placing said reservoir in fluid communication with the at least one second port of the handheld device.

17. The method of claim 12, wherein the treatment media comprises at least one of water and saline.

18. The method of claim 12, wherein the treatment media comprises a chemical agent.

19. The method of claim 12, wherein the chemical agent comprises at least one acid.

20. The method of claim 19, wherein the at least one acid comprises at least one of trichloroacetic acid (TCA), glycolic acid, alphahydroxy acid (AHA), lactic acid, citric acid and phenol.

* * * * *