



US005587497A

United States Patent [19]

DeLuca et al.

[11] **Patent Number:** 5,587,497[45] **Date of Patent:** Dec. 24, 1996[54] **19-NOR-VITAMIN D COMPOUNDS**

[75] Inventors: **Hector F. DeLuca**, Deerfield; **Heinrich K. Schnoes**; **Kato L. Perlman**, both of Madison, all of Wis.; **Rafal R. Scinski**, Warsaw, Poland; **Jean M. Prahl**, Madison, Wis.

[73] Assignee: **Wisconsin Alumni Research Foundation**, Madison, Wis.

[21] Appl. No.: 442,492

[22] Filed: **May 16, 1995**

Related U.S. Application Data

[62] Division of Ser. No. 281,261, Jul. 27, 1994, abandoned, which is a division of Ser. No. 123,485, Sep. 17, 1993, Pat. No. 5,342,975, which is a division of Ser. No. 960,241, Oct. 13, 1992, Pat. No. 5,246,925, which is a continuation of Ser. No. 879,706, May 5, 1992, abandoned, which is a continuation of Ser. No. 557,400, Jul. 3, 1990, abandoned, which is a division of Ser. No. 481,354, Feb. 16, 1990, Pat. No. 5,237,110, which is a continuation-in-part of Ser. No. 321,030, Mar. 9, 1989, abandoned.

[51] **Int. Cl.⁶** **C07C 401/00**

[52] **U.S. Cl.** **552/653**

[58] **Field of Search** 552/653; 514/167

[56] **References Cited****U.S. PATENT DOCUMENTS**

4,410,515	10/1983	Holick et al.	424/180
4,448,726	5/1984	DeLuca et al.	260/397
4,769,181	9/1988	DeLuca et al.	260/397
5,086,191	2/1992	DeLuca et al.	552/653
5,185,150	2/1993	DeLuca et al.	424/195
5,246,925	9/1993	DeLuca et al.	514/167
5,281,731	1/1994	DeLuca et al.	552/653
5,321,018	6/1994	DeLuca et al.	514/617
5,342,975	8/1994	DeLuca et al.	552/653
5,403,940	4/1995	Valle's et al.	514/167
5,428,029	6/1995	Doran et al.	514/167
5,449,668	9/1995	Sestelo et al.	514/167
5,451,574	9/1995	Baggiolini et al.	514/167

FOREIGN PATENT DOCUMENTS

0250755 1/1988 European Pat. Off. .
 WO85/03300 8/1985 WIPO .
 WO86/02649 5/1986 WIPO .

OTHER PUBLICATIONS

Chemical Abstracts (110:19004r) 1989.


Lythgoe et al "Calciferol And Its Relatives. Part 22. A Direct Total Synthesis of Vitamin D3", Journal of The Chemical Society, Perkin Transactions I, vol. 6, 1978, pp. 590-595.

Primary Examiner—Kimberly J. Prior

Attorney, Agent, or Firm—Andrus, Scales, Starke & Sawall

[57] **ABSTRACT**

This invention provides a novel class of vitamin D-related compounds, namely the 1 α -hydroxy-19-nor-vitamin D analogs, as well as a general method for their chemical synthesis. The compounds exhibit pronounced activity in arresting the proliferation of undifferentiated cells, including malignant cells, and in inducing their differentiation, and thus represent novel therapeutic agents for the treatment of malignant and other diseases characterized by the proliferative growth of undifferentiated cells. Formulations for therapeutic use and treatment methods are also provided. The 19-nor vitamin D compounds have the formula:



where X¹ and X² are each hydrogen or a hydroxy protecting group and R is a side chain.

12 Claims, No Drawings

19-NOR-VITAMIN D COMPOUNDS

This invention was made with United States government support awarded by the Department of Health and Human Services (NIH), Grant number: DK-14881. The United States Government has certain rights in this invention.

This application is a division of application Ser. No. 08/281,261 filed Jul. 27, 1994, now abandoned, which in turn is a divisional of Ser. No. 08/123,485 filed Sep. 17, 1993, now U.S. Pat. No. 5,342,975, which in turn is a divisional of Ser. No. 07/960,241 filed Oct. 13, 1992, now U.S. Pat. No. 5,246,925, which in turn is a continuation of Ser. No. 07/879,706 filed May 5, 1992, now abandoned, which in turn is a continuation of Ser. No. 07/557,400 filed Jul. 23, 1990, now abandoned, which in turn is a divisional of Ser. No. 07/481,354 filed Feb. 16, 1990, now U.S. Pat. No. 5,237,110, which in turn is a continuation-in-part application of Ser. No. 07/321,030 filed Mar. 9, 1989, now abandoned.


This invention relates to biologically active vitamin D compounds. More specifically, the invention relates to 19-nor-analogs of 1α -hydroxylated vitamin D compounds and to a general process for their preparation.

BACKGROUND


The 1α -hydroxylated metabolites of vitamin D—most importantly $1\alpha,25$ -dihydroxyvitamin D_3 and $1\alpha,25$ -dihydroxyvitamin D_2 —are known as highly potent regulators of calcium homeostasis in animals and humans, and more recently their activity in cellular differentiation has also been established. As a consequence, many structural analogs of these metabolites, such as compounds with different side chain structures, different hydroxylation patterns, or different stereochemistry, have been prepared and tested. Important examples of such analogs are 1α -hydroxyvitamin D_3 , 1α -hydroxyvitamin D_2 , various side chain fluorinated derivatives of $1\alpha,25$ -dihydroxyvitamin D_3 , and side chain homologated analogs. Several of these known compounds exhibit highly potent activity in vivo or in vitro, and possess advantageous activity profiles and thus are in use, or have been proposed for use, in the treatment of a variety of diseases such as renal osteodystrophy, vitamin D-resistant rickets, osteoporosis, psoriasis, and certain malignancies.

DISCLOSURE AND DESCRIPTION OF THE INVENTION

A class of 1α -hydroxylated vitamin D compounds not known heretofore are the 19-nor-analogs, i.e. compounds in which the ring A exocyclic methylene group (carbon 19) typical of all vitamin D system has been removed and replaced by two hydrogen atoms. Structurally these novel analogs are characterized by the general formula I shown below:




where X^1 and X^2 are each selected from the group consisting of hydrogen and acyl, and where the group R represents any of the typical side chains known for vitamin D type compounds. Thus, R may be an alkyl, hydroxyalkyl or fluoroalkyl group, or R may represent the following side chain:



wherein R^1 represents hydrogen, hydroxy or O-acyl, R^2 and R^3 are each selected from the group consisting of alkyl, hydroxyalkyl and fluoroalkyl, or, when taken together represent the group $-(CH_2)_m-$ where m is an integer having a value of from 2 to 5, R^4 is selected from the group consisting of hydrogen, hydroxy, fluorine, O-acyl, alkyl, hydroxyalkyl and fluoroalkyl, R^5 is selected from the group consisting of hydrogen, fluorine, alkyl hydroxyalkyl and fluoroalkyl, or, R^4 and R^5 taken together represent double-bonded oxygen, R^6 and R^7 are each selected from the group consisting of hydrogen, hydroxy, O-acyl, fluorine and alkyl, or, R^6 and R^7 taken together form a carbon-carbon double bond, and wherein n is an integer having a value of from 1 to 5, and wherein the carbon at any one of positions 20, 22, or 23 in the side chain may be replaced by an O, S, or N atom.

Specific important examples of side chains are the structures represented by formulas (a), (b), (c), (d) and (e) below, i.e. the side chain as it occurs in 25-hydroxyvitamin D_3 (a); vitamin D_3 (b); 25-hydroxyvitamin D_2 (c); vitamin D_2 (d); and the C-24-epimer of 25-hydroxyvitamin D_2 (e).




In this specification and the claims, the term 'alkyl' signifies an alkyl radical of 1 to 5 carbons in all isomeric forms, such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, etc., and the terms 'hydroxyalkyl' and 'fluoro-

3


alkyl' refer to such an alkyl radical substituted by one or more hydroxy or fluoro groups respectively, and the term 'acyl' means an aliphatic acyl group of 1 to 5 carbons, such as formyl, acetyl, propionyl, etc. or an aromatic acyl group such as benzoyl, nitrobenzoyl or halobenzoyl. The term 'aryl' signifies a phenyl-, or an alkyl-, nitro- or halo-substituted phenyl group.

The preparation of 1α -hydroxy-19-nor-vitamin D compounds having the basic structure shown above can be accomplished by a common general method, using known vitamin D compounds as starting materials. Suitable starting materials are, for example, the vitamin D compounds of the general structure II:



where R is any of the side chains as defined above. These vitamin D starting materials are known compounds, or compounds that can be prepared by known methods.

Using the procedure of DeLuca et al. (U.S. Pat. No. 4,195,027), the starting material is converted to the corresponding 1α -hydroxy-3,5-cyclovitamin D derivative, having the general structure III below, where X represents hydrogen and Q represents an alkyl, preferably methyl:





So as to preclude undesired reaction of the 1α -hydroxy group in subsequent steps, the hydroxy group is converted to the corresponding acyl derivative, i.e. the compound III shown above, where X represents an acyl group, using standard acylation procedures, such as treatment with an acyl anhydride or acyl halide- in pyridine at room temperature or slightly elevated temperature (30°-70° C.). It should be understood also that whereas the process of this invention is illustrated here with acyl protection of hydroxy functions, alternative standard hydroxy-protecting groups can also be used, such as, for example, alkylsilyl or alkoxyalkyl groups. Such protecting groups are well-known in the art. (e.g. trimethylsilyl, triethylsilyl, t.-butyldimethylsilyl, or tetrahy-

4

dروفuranyl methoxymethyl), and their use is considered a routine modification of experimental detail within the scope of the process of this invention.


The derivative as obtained above is then reacted with osmium tetroxide, to produce the 10,19-dihydroxy analog, IV (where X is acyl), which is subjected to diol cleavage using sodium metaperiodate or similar vicinal diol cleavage reagents (e.g. lead tetraacetate) to obtain the 10-oxo-intermediate, having the structure V below (where X is acyl):



These two consecutive steps can be carried out according to the procedures given by Paaren et al. [J. Org. Chem. 48, 3819 (1983)]. If the side chain unit, R, carries vicinal diols (e.g. 24,25-dihydroxy- or 25,26-dihydroxy, etc.), these, of course, also need to be protected, e.g. via acylation, silylation, or as the isopropylidene derivative prior to the periodate cleavage reactions.


In most cases, the acylation of the 1α -hydroxy group as mentioned above will simultaneously effect the acylation of side chain hydroxy functions, and these acylation conditions can, of course, be appropriately adjusted (e.g. elevated temperatures, longer reaction times) so as to assure complete protection of side chain vicinal diol groupings.

The next step of the process comprises the reduction of the 10-oxo-group to the corresponding 10-alcohol having the structure VI shown below (where X is acyl and Y represents hydroxy). When X is acyl, this reduction is carried out conveniently in an organic solvent at from about 0° C. to about room temperature, using NaBH_4 or equivalent hydride reducing agents, selective for the reduction of carbonyl groups without cleaving ester functions. Obviously, when X is a hydroxy-protecting group that is stable to reducing agents, any of the other hydride reducing agents (e.g. LiAlH_4 , or analogous reagents) may be employed also.



The 10-hydroxy intermediate is then treated with an alkyl- or arylsulfonylhalide (e.g. methanesulfonylchloride) in a suitable solvent (e.g. pyridine) to obtain the corresponding 10-O-alkyl- or arylsulfonyl derivative (the compound having the structure shown VI above where Y is alkyl-SO₂O-, or aryl-SO₂O-, and this sulfonate intermediate is then directly reduced, with lithium aluminum hydride, or the analogous known lithium aluminum alkyl hydride reagents in an ether solvent, at a temperature ranging from 0° C. to the boiling temperature of the solvent, thereby displacing the sulfonate group and obtaining the 10-deoxy derivative, represented by the structure VI above, where X and Y are both hydrogen. As shown by the above structure, a 1-O-acyl function in the precursor compound V is also cleaved in this reduction step to produce the free 1 α -hydroxy function, and any O-acyl protecting group in the side chain would, of course, likewise be reduced to the corresponding free alcohol function, as is well understood in the art. If desired, the hydroxy groups at C-1 (or hydroxy groups in the side chain) can be reprotected by acylation or silylation or ether formation to the corresponding acyl, alkylsilyl or alkoxyalkyl derivative, but such protection is not required. Alternative hydroxy-protecting groups, such as alkylsilyl or alkoxyalkyl groups would be retained in this reduction step, but can be removed, as desired, at this or later stages in the process by standard methods known in the art.

The above 1 α -hydroxy-10-deoxy cyclovitamin D intermediate is next solvolysed in the presence of a low-molecular weight organic acid, using the conditions of DeLuca et al. (U.S. Pat. Nos. 4,195,027 and 4,260,549). When the solvolysis is carried out in acetic acid, for example, there is obtained a mixture of 1 α -hydroxy-19-nor-vitamin D 3-acetate and 1 α -hydroxy-19-nor-vitamin D 1-acetate (compounds VII and VIII, below), and the analogous 1- and 3-acylates are produced, when alternative acids are used for solvolysis.



VI

5

10

15

20

25

30

35

40

45


50

55

60

6

-continued



VIII

Direct basic hydrolysis of this mixture under standard conditions then produces the desired 1 α -hydroxy-19-nor-vitamin D compounds of structure I above (where X¹ and X² are hydrogen). Alternatively, the above mixture of monoacetates may also be separated (e.g. by high pressure liquid chromatography) and the resulting 1-acetate and 3-acetate isomers may be subjected separately to hydrolysis to obtain the same final product from each, namely the 1 α -hydroxy-19-nor-vitamin D compounds of structure I. Also the separated monoacetates of structure VII or VIII or the free 1,3-dihydroxy compound can, of course, be reacylated according to standard procedures with any desired acyl group, so as to produce the product of structure I above where X¹ and X² represent acyl groups which may be the same or different.

Biological Activity of 1 α -Hydroxy-19-Nor-Vitamin D-Vitamin D Compounds

The novel compounds of this invention exhibit an unexpected pattern of biological activity, namely high potency in promoting the differentiation of malignant cells and little or no activity in calcifying bone tissue. This is illustrated by the biological assay results obtained for 1 α ,25-dihydroxy-19-nor-vitamin D₃ (compounds Ia), which are summarized in Tables 1 and 2, respectively. Table 1 shows a comparison of the activity of the known active metabolite 1 α ,25-dihydroxyvitamin D₃ and the 19-nor analog (Ia) in inducing the differentiation of human leukemia cells (HL-60 cells) in culture to normal cells (monocytes). Differentiation activity was assessed by three standard differentiation assays, abbreviated in Table 1 as NBT (nitroblue tetrazolium reduction), NSE (non-specific esterase activity), and PHAGO (phagocytosis activity). The assays were conducted according to known procedures, as given, for example, by DeLuca et al. (U.S. Pat. No. 4,717,721) and Ostrem et al., J. Biol. Chem. 262, 1416, 1987). For each assay, the differentiation activity of the test compounds is expressed in terms of the percent of HL-60 cells having differentiated to normal cells in response to a given concentration of test compound.

The results summarized in Table 1 clearly show that the new analog, 1 α ,25-dihydroxy-19-nor-vitamin D₃ (Ia) is as potent as 1 α ,25-dihydroxyvitamin D₃ in promoting the differentiation of leukemia cells. Thus in all three assays close to 90% of the cells are induced to differentiate by 1 α ,25-dihydroxy-vitamin D₃ at a concentration of 1 \times 10⁻⁷ molar, and the same degree of differentiation (i.e. 90, 84 and 90%) is achieved by the 19-nor analog (Ia).

TABLE 1

	Differentiation of HL-60 Cells		
	% Differentiated Cells (mean \pm SEM)		
	NBT	NSE	PRAGO
<u>1α,25-dihydroxyvitamin D₃ (moles/liter)</u>			
1 \times 10 ⁻⁷	86 \pm 2	89 \pm 1	87 \pm 3
1 \times 10 ⁻⁸	60 \pm 2	60 \pm 3	64 \pm 2
1 \times 10 ⁻⁹	33 \pm 2	31 \pm 2	34 \pm 1
<u>1α,25-Dihydroxy-19-nor- vitamin D₃, (Ia) (moles/liter)</u>			
2 \times 10 ⁻⁷	94 \pm 2	95 \pm 3	94 \pm 2
1 \times 10 ⁻⁷	90 \pm 4	84 \pm 4	90 \pm 4
5 \times 10 ⁻⁸	72 \pm 3	73 \pm 3	74 \pm 3
1 \times 10 ⁻⁸	61 \pm 3	60 \pm 3	56 \pm 1
1 \times 10 ⁻⁹	32 \pm 1	31 \pm 1	33 \pm 1

In contrast to the preceding results, the new 19-nor analog (Ia) exhibits no activity in an assay measuring the calcification of bone, a typical response elicited by vitamin D compounds. Relevant data, representing the results of an assay comparing the bone calcification activity in rats of 1 α ,25-dihydroxyvitamin D₃ and 1 α ,25-dihydroxy-19-nor-vitamin D₃ (Ia), are summarized in Table 2. This assay was conducted according to the procedure described by Tanaka et al., *Endocrinology* 92, 417 (1973).

The results presented in Table 2 show the expected bone calcification activity of 1 α ,25-dihydroxyvitamin D₃ as reflected by the increase in percent bone ash, and in total ash at all dose levels. In contrast, the 19-nor analog Ia exhibits no activity at all three dose levels, when compared to the vitamin D-deficient (-D) control group.

TABLE 2

Compound	Calcification Activity		
	Amount Administered* (pmoles/day/7 days)	% Ash (mean \pm SEM)	Total Ash (mg) (mean \pm SEM)
-D (control)	0	19 \pm 0.8	23 \pm 1.2
1 α ,25-dihydroxy- vitamin D ₃	32.5	23 \pm 0.5	34 \pm 1.6
	65.0	26 \pm 0.7	36 \pm 1.1
	325.0	28 \pm 0.9	40 \pm 1.9
1 α ,25-dihydroxy-19- nor-vitamin D ₃ (Ia)	32.5	22 \pm 0.9	28 \pm 1.6
	65.0	19 \pm 1.5	28 \pm 3.4
	325.0	19 \pm 1.2	30 \pm 2.4

*Each assay group comprised 6 rats, receiving the indicated amount of test compound by intraperitoneal injection daily for a period of seven days.

Thus the new 19-nor analog shows a selective activity profile combining high potency in inducing the differentiation of malignant cells with very low or no bone calcification activity. The compounds of this novel structural class, therefore, can be useful as therapeutic agents for the treatment of malignancies. Because the differentiative activity of vitamin D compounds on keratinocytes of skin (Smith et al., *J. Invest. Dermatol.* 86, 709, 1986; Smith et al., *J. Am. Acad. Dermatol.* 19, 516, 1988) is believed to be an indication of successful treatment of psoriasis (Takamoto et al., *Calc. Tissue Int.* 39, 360, 1986), these compounds should prove useful in treating this and other skin disorders characterized by proliferation of undifferentiated skin cells. These compounds should also find use in the suppression of parathy-

roid tissue, as for example, in cases of secondary hyperparathyroidism found in renal disease (Slatopolsky et al., *J. Clin. Invest.* 74, 2136, 1984).

For treatment purposes, the novel compounds of this invention can be formulated as solutions in innocuous solvents, or as emulsions, suspensions or dispersions in suitable innocuous solvents or carriers, or as pills, tablets or capsules, containing solid carriers according to conventional methods known in the art. For topical applications the compounds are advantageously formulated as creams or ointments or similar vehicle suitable for topical applications. Any such formulations may also contain other pharmaceutically-acceptable and non-toxic excipients such as stabilizers, anti-oxidants, binders, coloring agents or emulsifying or taste-modifying agents.

The compounds are advantageously administered by injection, or by intravenous infusion of suitable sterile solutions, or in the form of oral doses via the alimentary canal, or topically in the form of ointments, lotions, or in suitable transdermal patches. For the treatment of malignant diseases, the 19-nor-vitamin D compounds of this invention are administered to subjects in dosages sufficient to inhibit the proliferation of malignant cells and induce their differentiation into normal monocyte-macrophages. Similarly, for the treatment of psoriasis, the compounds may be administered orally or topically in amounts sufficient to arrest the proliferation of undifferentiated keratinocytes, and in the treatment of hyperparathyroidism, the compounds are administered in dosages sufficient to suppress parathyroid activity, so as to achieve parathyroid hormone levels in the normal range. Suitable dosage amounts are from 1 to 500 μ g of compound per day, such dosages being adjusted, depending on diseases to be treated, its severity and the response or condition of the subject as well-understood in the art.

This invention is more specifically described by the following illustrative examples. In these examples specific products identified by Roman numerals and letters, i.e. Ia, Ib, . . . , IIa, IIb, . . . , etc. refer to the specific structures and side chain combinations identified in the preceding description.

EXAMPLE 1

Preparation of 1 α ,25-dihydroxy-19-nor-vitamin D₃ (Ia)

(a) 1 α ,25-Dihydroxy-3,5-cyclovitamin D₃ 1-acetate, 6-methyl ether: Using 25-hydroxyvitamin D₃ (IIa) as starting material, the known 1 α ,25-dihydroxy-3,5-cyclovitamin D₃ derivative IIIa (X=H) was prepared according to published procedures (DeLuca et al., U.S. Pat. No. 4,195,027 and Paaren et al., *J. Org. Chem.* 45, 3252 (1980)). This product was then acetylated under standard conditions to obtain the corresponding 1-acetate derivative IIIa (X=Ac).

(b) 10,19-Dihydro-1 α ,25-tetrahydroxy-3,5-cyclovitamin D₃ 1-acetate, 6-methyl ether (IVa): Intermediate IIIa (X=Ac) was treated with a slight molar excess of osmium tetroxide in pyridine according to the general procedure described by Paaren et al. (*J. Org. Chem.* 48, 3819 (1983)) to obtain the 10,19-dihydroxylated derivative IVa. Mass spectrum m/z (relative intensity), 506 (M⁺, 1) 488 (2) 474 (40) 425 (45), 396 (15), 285 (5), 229 (30), 133 (45), 59 (80), 43 (100). ¹H NMR (CDCl₃) δ 0.5 (3H, s, 18-CH₃), 0.58 (1H, m, 3-H), 0.93 (3H, d, J=6.1 Hz, 21-CH₃), 1.22 (6H, s, 26-CH₃ and 27-CH₃), 2.10 (3H, s, COCH₃), 3.25 (3H, s, 6-OCR₃), 3.63 (2H, m, 19-CH₂), 4.60 (1H, d, J=9.2 Hz, 6-H), 4.63 (1H, dd, 1 β -H), 4.78 (1H, d, J=9.2 Hz, 7-H).

(c) 1 α ,25-Dihydroxy-10-oxo-3,5-cyclo-19-nor-vitamin D₃ 1-acetate, 6-methyl ether (Va): The 10,19-dihydroxylated

intermediate IVa was treated with a solution of sodium metaperiodate according to the procedure given by Paaren et al. (J. Org. Chem. 48, 3819, 1983) to produce the 10-oxo-cyclovitamin D derivative (Va, X=Ac). Mass spectrum m/z (relative intensity) 442 (M⁺-MeOH) (18), 424 (8), 382 (15), 364 (35), 253 (55), 225 (25), 197 (53), 155 (85), 137 (100). ¹H NMR (CDCl₃) δ 0.58 (3H, s, 18-CH₃), 0.93 (3H, d, J=6.6 Hz, 21-CH₃), 1.22 (6H, s, 26-CH₃ and 27-CH₃), 2.15 (s, 3-OCOCH₃), 3.30 (3H, s, 6-OCH₃), 4.61 (1H, d, J=9.1 Hz, 6-H), 4.71 (1H, d, J=9.6 Hz, 7-H), 5.18 (1H, m, 1β-H).

It has been found also that this diol cleavage reaction does not require elevated temperatures, and it is, indeed, generally preferable to conduct the reaction at approximately room temperature.

(d) 1α-Acetoxy-10,25-dihydroxy-3,5-cyclo-19-nor-vitamin D₃ 6-methyl ether (VIa, X=Ac, Y=OH): The 10-oxo derivative Va (X=Ac) (2.2 mg, 4.6 μmol) was dissolved in 0.5 ml of ethanol and to this solution 50 μl (5.3 μmol) of a NaBH₄ solution (prepared from 20 mg of NaBH₄, 4.5 ml water and 0.5 ml of 0.0 N NaOH solution) was added and the mixture stirred at 0° C. ca. 1.5 h, and then kept at 0° C. for 16 h. To the mixture ether was added and the organic phase washed with brine, dried over MgSO₄, filtered and evaporated. The crude product was purified by column chromatography on a 15x1 cm silica gel column and the alcohol VIa (X=Ac, Y=OH) was eluted with ethyl acetate hexane mixtures to give 1.4 mg (3 μmol) of product. Mass spectrum m/z (relative intensity) 476 (M⁺) (1), 444 (85), 426 (18), 384 (30), 366 (48), 351 (21), 255 (35), 237 (48), 199 (100), 139 (51), 59 (58).

(e) 1α,25-Dihydroxy-19-nor-vitamin D₃ (Ia, X¹=X²=H): The 10-alcohol (VIa, X=Ac, Y=OH) (1.4 mg) was dissolved in 100 μl anhydrous CH₂Cl₂ and 10 μl (14 μmol) triethylamine solution [prepared from 12 mg (16 μl) triethylamine in 100 μl anhydrous CH₂Cl₂], followed by 7 μl (5.6 μmol) mesyl chloride solution (9 mg mesyl chloride, 6.1 μl, in 100 μl anhydrous CH₂Cl₂) added at 0° C. The mixture was stirred at 0° C. for 2 h. The solvents were removed with a stream of argon and the residue (comprising compound VIa, X=Ac, Y=CH₃SO₂O-) dissolved in 0.5 ml of anhydrous tetrahydrofuran; 5 mg of LiAlH₄ was added at 0° C. and the mixture kept at 0° C. for 16 h. Excess LiAlH₄ was decomposed with wet ether, the ether phase was washed with water and dried over MgSO₄, filtered and evaporated to give the 19-nor product VIa (X=Y=H).

This product was dissolved in 0.5 ml of acetic acid and stirred at 55° C. for 20 min. The mixture was cooled, ice water added and extracted with ether. The ether phase was washed with cold 10% sodium bicarbonate solution, brine, dried over MgSO₄, filtered and evaporated to give the expected mixture of 3-acetoxy-1α-hydroxy- and 1α-acetoxy-3-hydroxy isomers, which were separated and purified by HPLC (Zorbax Sil column, 6.4x25 cm, 2-propanol in hexane) to give about 70 μg each of compounds VIIa and VIIIa. UV (in EtOH) λ_{max} 242.5 (OD 0.72), 251.5 (OD 0.86), 260 (OD 0.57).

Both 19-nor-1,25-dihydroxyvitamin D₃ acetates VIIa and VIIIa were hydrolyzed in the same manner. Each of the monoacetates was dissolved in 0.5 ml of ether and 0.5 ml 0.1 N KOH in methanol was added. The mixture was stirred under argon atmosphere for 2 h. More ether was added and the organic phase washed with brine, dried over anhydrous MgSO₄, filtered and evaporated. The residue was dissolved in a 1:1 mixture of 2-propanol and hexane and passed through a Sep Pak column and washed with the same solvent. The solvents were evaporated and the residue purified by HPLC (Zorbax Sil, 6.4x25 cm, 10% 2-propanol in hexane). The hydrolysis products of VIIa and VIIIa were identical and gave 66 μg of Ia (X¹=X²=H). Mass spectrum

(m/z relative intensity) 404 (M⁺) (100), 386 (41), 371 (20), 275 (53), 245 (51), 180 (43), 135 (72), 133 (72), 95 (82), 59 (18), exact mass calcd. for C₂₆H₄₄O₃ 404.3290, found 404.3272. ¹H NMR (CDCl₃) δ 0.52 (3H, s, 18-CH₃), 0.92 (3H, d, J=6.9 Hz, 21-CH₃), 1.21 (6H, s, 26-CH₃ and 27-CH₃), 4.02 (1H, m, 3α-H), 4.06 (1H, m, 1β-H), 5.83 (1H, d, J=11.6 Hz, 7-H), 6.29 (1H, d, J=10.7 Hz, 6-H). UV (in EtOH) λ_{max} 243 (OD 0.725), 251.5 (OD 0.823), 261 (OD 0.598).

EXAMPLE 2

Preparation of 1α-hydroxy-19-nor-vitamin D₃ (Ib)

(a) With vitamin D₃ (IIb) as starting material, and utilizing the conditions of Example 1a, there is obtained known 1α-hydroxy-3,5-cyclovitamin D₃ 1-acetate, 6-methyl ether, compound IIIb (X=Ac).

(b) By subjecting intermediate IIIb (X=Ac), as obtained in Example 2a above to the conditions of Example 1b, there is obtained 10,19-dihydro-1α,10,19-trihydroxy-3,5-cyclovitamin D₃ 1-acetate, 6-methyl ether IVb (X=Ac).

(c) By treatment of intermediate IVb (X=Ac) with sodium metaperiodate according to Example 1c above, there is obtained 1α-hydroxy-10-oxo-3,5-cyclo-19-nor-vitamin D₃ 1-acetate, 6-methyl ether Vb (X=Ac).

(d) Upon reduction of the 10-oxo-intermediate Vb (X=Ac) under the conditions of Example 1d above, there is obtained 1α-acetoxy-10-hydroxy-3,5-cyclo-19-nor-vitamin D₃ 6-methyl ether VIb (X=Ac, Y=OH).

(e) Upon processing intermediate VIb (X=Ac, Y=OH) through the procedure given in Example 1e above, there is obtained 1α-hydroxy-19-nor-vitamin D₃ (Ib, X¹=X²=H).

EXAMPLE 3

Preparation of 1α,25-dihydroxy-19-nor-vitamin D₂

(a) Utilizing 25-hydroxyvitamin D₂ (IIc) as starting material and experimental conditions analogous to those of Example 1a, there is obtained 1α,25-dihydroxy-3,5-cyclovitamin D₂ 1-acetate, 6-methyl ether, compound IIIc (X=Ac).

(b) Subjecting intermediate IIIc (X=Ac), as obtained in Example 3a above, to the reaction conditions of Example 1b, provides 10,19-dihydro-1α,10,19,25-tetrahydroxy-3,5-cyclovitamin D₂ 1-acetate, 6-methyl ether, IVc (X=Ac).

(c) By treatment of intermediate IVc (X=Ac) with sodium metaperiodate according to general procedures of Example 1c above, there is obtained 1α,25-dihydroxy-10-oxo-3,5-cyclo-19-nor-vitamin D₂ 1-acetate, 6-methyl ether Vc (X=Ac).

(d) Upon reduction of the 10-oxo-intermediate Vc (X=Ac) under conditions analogous to those of Example 1d above, there is obtained 1α-acetoxy-10,25-dihydroxy-3,5-cyclo-19-nor-vitamin D₂ 6-methyl ether VIc (X=Ac, Y=OH).

(e) Upon processing intermediate VIc (X=Ac, Y=OH) through the procedural steps given in Example 1e above, there is obtained 1α,25-dihydroxy-19-nor-vitamin D₂ (Ic, X¹=X²=H).

EXAMPLE 4

Preparation of 1α-hydroxy-19-nor-vitamin D₂

(a) With vitamin D₂ (IIc) as starting material, and utilizing the conditions of Example 1a, there is obtained known 1α-hydroxy-3,5-cyclovitamin D₂ 1-acetate, 6-methyl ether, compound IIIc (X=Ac).

11

(b) By subjecting intermediate IIIId (X=Ac), as obtained in Example 4a above to the conditions of Example 1b, there is obtained 10,19-dihydro-1 α ,10,19-trihydroxy-3,5-cyclo-vitamin D₂ 1-acetate, 6-methyl ether, IVd (X=Ac).


(c) By treatment of intermediate IVb (X=Ac) with sodium metaperiodate according to Example 1c above, there is obtained 1 α -hydroxy-10-oxo-3,5-cyclo-19-nor-vitamin D₂ 1-acetate, 6-methyl ether, Vd (X=Ac).

(d) Upon reduction of the 10-oxo-intermediate Vd (X=Ac) under the conditions of Example 1d above, there is obtained 1 α -acetoxy-10-hydroxy-3,5-cyclo-19-nor-vitamin D₂ 6-methyl ether, VIId (X=Ac, Y=OH).


(e) Upon processing intermediate VIId (X=Ac, Y=OH) through the procedure given in Example 1e above, there is obtained 1 α -hydroxy-19-nor-vitamin D₂ (Id, X¹=X²=H).

We claim:

1. A compound having the formula



where X¹ and X² are each selected from hydrogen, acyl, alkylsilyl and alkoxyalkyl, and where R is selected from alkyl, hydrogen, hydroxyalkyl, said hydroxyalkyl having from 4 to 5 carbon atoms, fluoroalkyl and a side chain of the formula




wherein R¹ represents hydrogen, hydroxy or O-acyl, R² and R³ are each selected from alkyl, hydroxyalkyl and fluoroalkyl, or, when taken together represent the group $-(CH_2)_m-$ where m is an integer having a value of from 2 to 5, R⁴ is selected from hydrogen, hydroxy, fluorine, O-acyl, alkyl, hydroxyalkyl and fluoroalkyl, R⁵ is selected from hydrogen, fluorine, alkyl, hydroxyalkyl and fluoroalkyl, or R⁴ and R⁵ taken together represent double-bonded oxygen, R⁶ and R⁷ are each selected from hydrogen, hydroxy, O-acyl, fluorine and alkyl, or, R⁶ and R⁷ taken

12


together form a carbon-carbon double bond, and wherein n is an integer having a value of from 1 to 5 and wherein any of the groups $-CH(CH_3)-$, $-CH(R^7)-$, or $-CH(R^6)-$ at positions 20, 22 and 23, respectively, may be replaced by an oxygen atom, with the proviso that when n is 2 to 5 each R⁴ is independently selected from hydrogen, hydroxy, fluorine, O-acyl, alkyl, hydroxyalkyl and fluoroalkyl, and each R⁵ is independently selected from hydrogen, fluorine, alkyl, hydroxyalkyl and fluoroalkyl.

2. A compound according to claim 1 where X¹ and X² represent hydrogen, and where R¹ is hydroxy, both of R² and R³ are selected from the group consisting of methyl, trifluoromethyl, ethyl and propyl, both of R⁶ and R⁷ are hydrogen, or together form a carbon-carbon double bond, R⁴ and R⁵ are hydrogen and n is an integer having the values 1, 2 or 3.


3. The compound of claim 1 where R is a side chain of the formula




4. The compound of claim 1 where R is a side chain of the formula



5. The compound of claim 1 where R is a side chain of the formula



6. The compound of claim 1 where R is a side chain of the formula



7. 1 α ,25-dihydroxy-19-nor-vitamin D₃.

8. 1 α -hydroxy-19-nor-vitamin D₃.

9. 1 α ,25-dihydroxy-19-nor-vitamin D₂.

10. 1 α -hydroxy-19-nor-vitamin D₂.

11. 1 α -hydroxy-19-nor-24 epi-vitamin D₂.

12. 1 α ,25-dihydroxy-19-nor-24 epi-vitamin D₂.

* * * * *